

LETTERS

J. Raptor Res. 53(2):220–222
© 2019 The Raptor Research Foundation, Inc.

PREDATION OF SOUTHERN RED OCTOPUS (*ENTEROCTOPUS MEGALOCYATHUS*) BY STRIATED CARACARAS (*PHALCOBOENUS AUSTRALIS*) IN THE FALKLAND ISLANDS

KATIE J. HARRINGTON¹ AND KEITH L. BILDSTEIN

Acopian Center for Conservation Learning, Hawk Mountain Sanctuary, 410 Summer Valley Road, Orwigsburg, PA 17961 USA

KEY WORDS: *Striated Caracara; Phalcoboenus australis; southern red octopus; Enteroctopus megalocyathus; Cephalopod; diet; dietary flexibility; Falconid; island; Islas Malvinas; raptor.*

Globally near-threatened Striated Caracaras (*Phalcoboenus australis*; Birdlife International 2017) are scavenging and occasionally predatory birds of prey (Falconiformes) restricted to exposed islands in southernmost South America and the Falkland Islands (Isla Malvinas). During the austral summer, this inquisitive and gregarious species feeds primarily on seabird eggs, nestlings, and dead and dying adult seabirds at seabird colonies (Strange 1996, Catry et al. 2008), supplemented by marine and terrestrial invertebrates, e.g., Gastropoda and Diptera (Strange 1996, Rexer-Huber and Bildstein 2013), and placentas and carrion of marine mammals and livestock (Strange 1996, Catry et al. 2008). Juvenile caracaras often vie for carcasses in “gangs” of up to several dozen birds (Strange 1996, Catry et al. 2008, Rexer-Huber and Bildstein 2013), not unlike juvenile Common Ravens (*Corvus corax*) in the northern hemisphere (Heinrich and Marzluff 1991). In the winter months, when seabirds are largely foraging at sea, Saunders Island caracaras become food-stressed and lose body mass (Rexer-Huber and Bildstein 2013), and invertebrates and carrion may become more important resources, allowing them to subsist until the seabirds return.

The southern red octopus (*Enteroctopus megalocyathus*) is a commercially fished large benthic species with a sub-Antarctic distribution (Ortiz et al. 2011) found from the intertidal zone to a depth of roughly 140 m (Ibáñez and Chong 2008). To our knowledge, the only published reports of southern red octopus predators within the waters surrounding the Falkland Islands include southern sea lions (*Otaria flavescens*; Shallow Marine Surveys Group 2010), Gentoo Penguins (*Pygoscelis papua*; Clausen et al.

2005), and spiny dogfish (*Squalus acanthias*; Laptikhovsky et al. 2001).

Saunders Island is a “nursery island” in the Falklands archipelago that supports over 100 Striated Caracaras throughout the year, most of which are nonbreeding juveniles and subadults (Harrington et al. 2018). On 10 February 2017, we observed 31 caracaras prey upon a southern red octopus (est. 3–4 kg in mass) in the exposed rocky intertidal area adjacent to a south-facing sandy beach on Saunders Island, Falkland Islands (51°18.59'S, 60°14.32'W). Prior to the predation event, the foraging group included 12 juveniles, three subadults, and one adult (aged according to plumage characteristics described by Strange 1996) that appeared to be searching for gastropods and other prey made available by a low spring tide. The birds pulled limpets (*Fissurella picta* and *Patinigera* spp.) from exposed overhanging rocks, crevices, and kelp (*Lessonia* spp.), and additional caracaras arrived to the 50-m² rocky intertidal zone as the tide receded.

At 1455 H (GMT–3 hr), a juvenile caracara found the octopus in a shallow tidal pool (<20 cm deep). The caracara took hold of one of the octopus' arms in its bill and pulled it to full extension. Two juvenile caracaras foraging within 1 m of the bird that first struck the octopus then joined in preying on the octopus, and one of the caracaras dismembered an arm by holding down the arm with its foot, seizing it in its bill, and tearing it away. The birds dislodged the octopus and pulled it onto the surface of an exposed rock where, inverted, the octopus tried to right itself (Fig. 1). Within 1 min of discovery, nine juvenile and three subadult caracaras had encircled and attacked the octopus. Five juveniles and two subadults bit off chunks of arm and retreated several meters to consume them. Within 2 min, an additional nine juveniles, eight subadults, and three adults had arrived. In total, 31 caracaras encircled the octopus. At 1457 H, the octopus still had three arms intact, as well as its head and mantle. By 1501 H, only the head and mantle remained, which the caracaras were able to grab with their talons and carry for 2–5 m,

¹ Present address: Moss Landing Marine Laboratories, 8272 Moss Landing Road, Moss Landing, CA 95039 USA; email address: katiejharrington@gmail.com



Figure 1. Juvenile and subadult Striated Caracaras encircling and consuming southern red octopus, 10 February 2017. Photo by K. Harrington.

before consuming them. The feeding event lasted less than 8 min.

Striated Caracaras complement their mainly seabird and insectivorous diet with a variety of other food resources (Strange 1996, Rexer-Huber and Bildstein 2013). Pellet analyses show squid remains in the diet of caracaras on New Island, 90 km to the west-southwest of Saunders Island (Catry et al. 2008), likely because Striated Caracaras consume spilled regurgitated food brought ashore by penguins for their young (Strange 1996, K. Harrington unpubl. data). Juvenile and adult Striated Caracaras have long been noted for experimenting with and testing their environment to determine the availability of suitable food resources (Darwin 1845, Strange 1996, Rexer-Huber et al. 2012). Our observation further highlights the species' dietary flexibility, a characteristic that may provide resilience in the face of environmental change and an adaptation to seasonal variations in food abundance in their island habitats. To our knowledge, there is no other record of a Falconid predating an octopus.

Funding for this study was provided by Wallace Research Foundation, Chris and Stanley Clayton, Jim and Jean Macaleer, and Hawk Mountain Sanctuary. We thank the Falkland Islands Government Environmental

Planning Department for providing permission to work on the islands and the Pole-Evans family for being gracious hosts and providing logistical support during our stays on Saunders Island. We give further thanks to Dr. Louise Allcock, National University of Ireland, Galway, for identifying the octopus and to Jonathan Meiburg and two anonymous reviewers who provided critical comments that improved an earlier version of this letter. All research was conducted in accordance with the Falkland Island Government Conservation of Wildlife and Nature Ordinance under Research License No. R22/2015. This is Hawk Mountain Contribution to Conservation Science No. 310.

LITERATURE CITED

- BirdLife International (2017). Species factsheet: *Phalcoboenus australis*. <http://www.birdlife.org>.
- Catry, P., M. Lecoq, and I. Strange (2008). Population growth and density, diet and breeding success of Striated Caracaras *Phalcoboenus australis* on New Island, Falkland Islands. *Polar Biology* 31:1167–1174.
- Clausen, A., A. Arkhipkin, V. Laptikhovsky, and N. Huin (2005). What is out there: diversity in feeding of Gentoo Penguins (*Pygoscelis papua*) around the Falkland Islands (southwest Atlantic). *Polar Biology* 28:653–662.

- Darwin, C. R. (1845). *Journal of Researches into the Natural History and Geology of Countries Visited During the Voyage of HMS Beagle Round the World*. John Murray, London, UK.
- Harrington, K. J., S. Pole-Evans, M. Reeves, M. Bechard, M. Bobowski, D. R. Barber, K. Rexer-Huber, N. Lecomte, and K. L. Bildstein (2018). Seasonal micro-migration in a farm-island population of Striated Caracaras (*Phalcoboenus australis*) in the Falkland Islands. *Movement Ecology* 6:4. doi:10.1186/s40462-018-0122-8.
- Heinrich, B., and J. M. Marzluff (1991). Do Common Ravens yell because they want to attract others? *Behavioral Ecology and Sociobiology* 28:13–21.
- Ibáñez, C. M., and J. V. Chong (2008). Feeding ecology of *Enteroctopus megalocyathus* (Cephalopoda: Octopodidae) in southern Chile. *Journal of the Marine Biological Association of the United Kingdom* 88:793–798.
- Laptikhovsky, V. V., A. I. Arkhipkin, and A. C. Henderson (2001). Feeding habits and dietary overlap in spiny dogfish *Squalus acanthias* (Squalidae) and narrowmouth catshark *Schroederichthys bivius* (Scyliorhinidae). *Journal of the Marine Biological Association of the United Kingdom* 81:1015–1018.
- Ortiz, N., M. E. Ré, F. Márquez, and N. G. Glembocki (2011). The reproductive cycle of the red octopus *Enteroctopus megalocyathus* in fishing areas of northern Patagonian coast. *Fisheries Research* 110:217–223.
- Rexer-Huber, K., and K. L. Bildstein (2013). Winter diet of Striated Caracara *Phalcoboenus australis* (Aves, Polyborinae) at a farm settlement on the Falkland Islands. *Polar Biology* 36:437–443.
- Rexer-Huber, K., G. C. Parker, M. Reeves, A. Stanworth, and R. J. Cuthbert (2012). Baiting Trials, Winter Biology and Non-target Species: House Mice on Steeple Jason August–September 2012. RSPB Research Report No. 51. Royal Society for the Protection of Birds, Sandy, Bedfordshire, UK.
- Shallow Marine Surveys Group (2010). *Marine Resources of the Falklands' Shallow Marine Environment: A Review of Species, Distributions, and Sustainability*. Report of the Falkland Islands Government, Stanley, Falkland Islands.
- Strange, I. (1996). The Striated Caracara *Phalcoboenus australis* in the Falkland Islands. Philip Myers, Warrington, UK.

Received 17 August 2018; accepted 18 December 2018
Associate Editor: Christopher W. Briggs